

Experimental Basosquamous Carcinoma Model in Rats

Snarskaya E., Pylev L., Akhunzyanov A., Kuznetcova E.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Previous studies have established that 7,12-dimethylbenz(a)anthracene (DMBA) can initiate skin tumorigenesis in conventional furred mouse models by acting on hair follicle stem cells. In this work, we have developed a simple and convenient rat model of basosquamous carcinoma (BSC) using DMBA-induced carcinogenesis in male specific pathogen free (SPF) Wistar rats with no additional tumor promoter agents. The results showed that topical application of 0.1% solution of DMBA in benzene in a volume of 40 µL twice a week produced skin tumors after 8–9 months. As a result, during the 11–12th months, we obtained 15 animals with BSCs, 22 with basal cell carcinomas (BCCs), and 3 with squamous cell carcinomas (SCCs). This chemically driven skin cancer model in Wistar rats can serve as a suitable alternative to the mouse skin cancer model and can be reliably replicated as demonstrated by the experiment.

<http://dx.doi.org/10.1007/s12668-016-0380-0>

Keywords

Basosquamous carcinoma, Chemical carcinogenesis, DMBA, Rat model

References

- [1] Neagu, M., et al. (2016). Chemically induced skin carcinogenesis: updates in experimental models (review). *Oncol Rep*, 35, 2516–2528.
- [2] DiGiovanni, J. (1992). Multistage carcinogenesis in mouse skin. *Pharmacol Ther*, 54, 63–128.
- [3] Abel, E., Angel, J., Kiguchi, K., & DiGiovanni, J. (2009). Multistage chemical carcinogenesis in mouse skin: fundamentals and applications. *Nat Protoc*, 4, 1350–1362.
- [4] Uhmman, A., et al. (2014). DMBA/TPA treatment is necessary for BCC formation from patched deficient epidermal cells in *Ptch(flox/flox)CD4Cre(+/-)* mice. *J Invest Dermatol.*, 134, 2620–2629.
- [5] Miller, S., Wei, Z., Wilson, C., Dzubow, L., Sun, T., & Lavker, R. (1993). Mouse skin is particularly susceptible to tumour initiation during early anagen of the hair cycle: possible involvement of hair follicle stem cells. *J Invest Dermatol*, 101, 591–594.
- [6] Iversen, O. H. (1994). A course of very small doses of DMBA, each of them allegedly with no promoting potency, acts with clear synergistic effect as a strong promoter of DMBA-initiated mouse skin carcinogenesis. A comparison of the tumourigenic and carcinogenic effects of DMBA (7,12-dimethylbenz- α -anthracene) and TPA (12-Otetradecanoyl-phorbol-13-acetate) used as initiators and promoters in classical two-stage experimental protocols. *APMIS Suppl*, 41, 1–38.
- [7] Mughal, M., Azizullah, & Ahmed, T. (2005). Carcinogenic effects of DMBA and TPA on skin of mice. *Biologia (Lahore)*, 51, 201–206.
- [8] Thomas, G., et al. (2016). Studying skin tumourigenesis and progression in immunocompetent hairless SKH1-hr mice using chronic 7,12-dimethylbenz(a)anthracene topical applications to develop a useful experimental skin cancer model. *Lab Anim*. doi:10.1177/0023677216637305.

- [9] Garcia, C., Poletti, E., & Crowson, A. (2009). Basosquamous carcinoma. *J Am Acad Dermatol*, 60, 137–143.
- [10] Braun-Falco, O., Plewig, G., & Wolf, W. H. C. (2000). *Dermatologica* (2nd ed.pp. 1481–1482). Berlin: Springer-Verlag.
- [11] Darier, J., & Ferrand, M. (1955). Zepitelioma ravimenteux mixte et intermediaire. *Ann Dermatol lyphiligr (Paris)*, 82, 12–14.
- [12] Lever W.F., Schaumburg-Lever G. (1983). *Histopathology of the skin*. 6th ed. Philadelphia.
- [13] Snarskaja, E. S. (2014). Modern and promising therapy of different forms of basal cell skin cancer and metatypical skin cancer. *Russian Journal of skin and sexually transmitted diseases.*, 1, 4–9.
- [14] Hlebnikova, A. N., & Kazanceva, I. A. (2000). Histological variants of basal cell skin cancer (10.4% of cases of metatypical cancer). *Russian Journal of skin and sexually transmitted diseases*, 2, 4–8.
- [15] Mancuso, M., et al. (2009). Modulation of basal and squamous cell carcinoma by endogenous estrogen in mouse models of skin cancer. *Carcinogenesis*, 30, 340–347.
- [16] Lu, S., Herrington, H., & Wang, X. (2006). Mouse models for human head and neck squamous cell carcinomas. *Head Neck*, 28, 945–954.
- [17] Björkhem-Bergman, L., Ekström, L., & Eriksson, L. (2012). Review: exploring anticarcinogenic agents in a rat hepatocarcinogenesis model—focus on selenium and statins. *In Vivo*, 26, 527–535.
- [18] Langdon, S. (2012). Animal modeling of cancer pathology and studying tumor response to therapy. *Curr Drug Targets*, 13, 1535–1547.
- [19] Perše, M., & Cerar, A. (2011). Morphological and molecular alterations in 1,2-dimethylhydrazine and azoxymethane induced colon carcinogenesis in rats. *J Biomed Biotechnol*, 2011, 473964.
- [20] McKiernan, M. V., & McCann, J. J. (1995). Collision tumour—an unusual skin lesion. *Br J Dermatol*, 132, 486.